Charmonia from SQM 2016




Songkyo Lee




dilepton meeting 6th July 2016




• J/ψ in PbPb @ 5 TeV - Paper submitted (http://arxiv.org/abs/1202.1383)




- finer binning than 2.76 TeV
- Similar trends (slightly higher)
 - 5.02 TeV $R_{AA}^{0-90\%}$ = 0.66 ± 0.01(stat) ± 0.05(syst)
 - 2.76 TeV $R_{AA}^{0-90\%}$ = 0.58 ± 0.01(stat) ± 0.09(syst)



• J/ψ in PbPb @ 5 TeV - Paper submitted (http://arxiv.org/abs/1202.1383)




- very low p_T J/ψ excess
- presumably originated from photon-production
- p_T > 0.3 GeV cut applied
 to reduce these contributions







- 5.02 TeV / 2.76 TeV
- Competition between suppression vs regeneration
- Regeneration more dominant?

$$r^{0-10\%} = 1.17 \pm 0.04(\text{stat}) \pm 0.20(\text{syst})$$



- Transport : rate equation of suppression and regeneration by/in the QGP
- Statistical hadronization : all J/ ψ produced by statistical hadronization at the QGP phase boundary
- Co-movers : suppression by the co-moving partonic medium + regeneration







- Hint for an increase with $\sqrt{s_{NN}}$ at $p_T = 2 6$ GeV
- Regeneration & radial flow effects?

Other LHC experiments




No new results from ATLAS and LHCb yet







- LHCb also working on..
 - photo-production
 - fixed target p-[He,Ne,Ar] e.g. @ 110 GeV



PHENIX




Double ratio [ψ(2S) / J/ψ] in p-Al @ 200 GeV




- Hint for stronger suppression at backward
- Co-mover model qualitatively agrees with data

STAR








- STAR: new, independent measurement at |y|<0.5 via muon channel
- Confirms existing picture:
 - ▶ Low-p_T J/ψ at the LHC are less suppressed than at RHIC, extra source
 - ▶ High-p_T J/ψ at the LHC are more suppressed than at RHIC, more dissociation


Transport approch




Xiaojian Due, Tuesday



- ψ(2S) regenerated later than J/ψ
- flow pushes $\psi(2S)$ to higher p_T






plan to provide prediction for our 5.02 TeV results


(~ before the end of summer)


back up

10



- Very-low-p_T J/ψ excess
 - Seen in peripheral Pb-Pb collisions at 2.76 TeV
 - Presumably of EM origin






- Due to the very specific origin and kinematics, photo-produced J/ψ could become an useful probe of the QGP
- In the mean time, it constitutes a "contamination" to the hadronic R_{AA}
 - Apply a cut $p_T > 0.3$ GeV to reduce photo-production contribution


PHENIX




Backward: Al-going direction




central collision




Forward: p-going direction



Backward: Au-going direction




central collision




Forward: p-going direction


Run-15 p+Au √s = 200 GeV





STAR
















- Coherent and incoherent photoproduction of ρ mesons observed in Ultra-Peripheral Collisions (UPC)
- Observe excess of very low p_T J/ ψ in peripheral collisions with features consistent with coherent photoproduction
 - Similar slope as UPC: 199 ±31(GeV/c)⁻²
 UPC in STARLIGHT: 196 (GeV/c)⁻²


interactions


STAR














- Coherent and incoherent photoproduction of ρ mesons observed in Ultra-Peripheral Collisions (UPC)
- Observe excess of very low p_T J/ψ in peripheral collisions with features consistent with coherent photoproduction
 - Similar slope as UPC: 199 ±31(GeV/c)⁻²
 UPC in STARLIGHT: 196 (GeV/c)⁻²
 - Production cross-section independent of centrality

A challenge for theory but a new opportunity for QGP studies?

7honus Vo

Photon-nucleus

interactions