

Belle Analysis Status

18th, April, 2018

Seongbae Yang
Department of Physics
Korea University

Events Number Test


- Data sample for this test

→ Only Exp65 ($\sim 49.7 \text{ fb}^{-1}$, $\sim 5\%$ of total data sample)

- Λ Reconstruction

→ Used "Mdst_vee2".

2 and 3 in the vee2 collection




*Lambda cut condition: 'GoodLambda 1 or 2' and ' $M(p\pi^-)$ in $\sim \pm 5\sigma$ '
 → Mass constraint fit applied.

■ Λ_c^+ Reconstruction

→ Adding one pi+ to the Lambda

pi+ cut condition: $R(\pi|p) > 0.4$, $R(\pi|K) > 0.4$, $R(e) < 0.9$, $|ip_r| < 0.1$ cm,
 $|ip_z| < 2$ cm, and more than 1 hit for both SVD r-phi and z.

→ Λ_c^+ cut condition: $x_p > 0.5$




→ Expected yield in total data sample: ~230000 events.

■ New results from Belle


#BN1467, $\Omega^{*-} \rightarrow \Xi^0 K^-$ & $\Xi^- K_s^0$

Clear peak at ~ 2012 MeV/ c^2





■ Events number test

- Data sample: $\Upsilon(1s)$ resonance, $\sim 5.7 \text{ fb}^{-1}$ ($\sim 17\%$ of $\Upsilon(1s, 2s, 3s)$)
- Same Lambda selection with Λ_c^+



→ Adding one K- to the Lambda

π^+ cut condition: $R(K|\pi) > 0.9$, $R(K|p) > 0.4$, $R(e) < 0.9$, $|ip_r| < 20.0 \text{ cm}$, $|ip_z| < 20.0 \text{ cm}$, and more than 1 hit for both SVD r-phi and z.

$\gamma(1s)$ resonance $\gamma(1s)$ off-resonance

→ $\Omega^- \rightarrow \Lambda K^-$: 60000 events in $\gamma(1s, 2s, 3s)$ data.